



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A01N 63/00, 57/00, 25/00	A1	(11) International Publication Number: WO 00/41567 (43) International Publication Date: 20 July 2000 (20.07.00)												
(21) International Application Number: PCT/US00/00954 (22) International Filing Date: 14 January 2000 (14.01.00) (30) Priority Data: <table border="0"><tr><td>60/116,085</td><td>15 January 1999 (15.01.99)</td><td>US</td></tr><tr><td>60/117,559</td><td>28 January 1999 (28.01.99)</td><td>US</td></tr><tr><td>60/131,662</td><td>29 April 1999 (29.04.99)</td><td>US</td></tr><tr><td>09/448,245</td><td>24 November 1999 (24.11.99)</td><td>US</td></tr></table> (71) Applicant: COGNIS CORPORATION [US/US]; Suite 200, 2500 Renaissance Boulevard, Gulph Mills, PA 19406 (US). (72) Inventor: BRIGANCE, Mickey, R.; Suite #6, 1385 West Brierbrook, Germantown, TN 38138 (US). (74) Agent: DRACH, John E.; Cognis Corporation, Suite 200, 2500 Renaissance Boulevard, Gulph Mills, PA 19406 (US).		60/116,085	15 January 1999 (15.01.99)	US	60/117,559	28 January 1999 (28.01.99)	US	60/131,662	29 April 1999 (29.04.99)	US	09/448,245	24 November 1999 (24.11.99)	US	(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
60/116,085	15 January 1999 (15.01.99)	US												
60/117,559	28 January 1999 (28.01.99)	US												
60/131,662	29 April 1999 (29.04.99)	US												
09/448,245	24 November 1999 (24.11.99)	US												
(54) Title: ADJUVANT COMPOSITION (57) Abstract <p>An adjuvant composition for pesticide formulations which exhibits reduced eye irritancy is comprised of a polyoxyalkylene aliphatic amine containing at least about 2 moles of an alkylene oxide group and an eye irritation reducing compound which simultaneously reduces the eye irritation caused by the polyoxyalkylene aliphatic amine and which complexes with metal ions such as calcium and iron which may be present in an aqueous solution of the adjuvant and a pesticide or a plant growth regulator. The eye irritation reducing compound is a carboxylic acid that is capable of complexing with or forming chelates with metal ions in aqueous solution while simultaneously reducing the eye irritation caused by the polyoxyalkylene aliphatic amine. The adjuvant also contains a mixture of polyhydric alcohols and, optionally, a defoamer. The adjuvant is particularly in N-phosphonomethylglycine (glyphosate) herbicidal formulations.</p>														

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

ADJUVANT COMPOSITION

BACKGROUND OF THE INVENTION

5 Insecticides, insect repellents, fungicides, bactericides, bacteriostats, herbicides, and plant growth regulators are normally formulated into various products for use on crops, for insect control, weed control and the like. Alternatively, the products may be formulated as liquids or powders or granules. Solvents, emulsifiers, dispersing agents and wetting agents are normally incorporated into such compositions to ensure the preparation of a uniform
10 pesticide formulation.

These formulation components are also selected to ensure that the pesticide composition will disperse or emulsify evenly in a tank mix at the point of application. They also have a third purpose that is to ensure optimum delivery of the tank mix preparation to the targeted pest or substrate. Sometimes the
15 surfactants incorporated in pesticide formulations are not sufficient to fully ensure stable tank mixes when such tank mixes contain multiple components. Similarly, it may be necessary to add adjuvants to the tank mix for full stability. Chemically, surfactants are the most important and widely used adjuvants. Surfactants may affect many properties of the formulation such as solubility, volatility, specific
20 gravity, corrosiveness, efficacy, and freezing and flash points. It is widely known that adding surfactant-based adjuvants to the tank mix will realize the desired stabilization. Adjuvants also potentiate pesticidal activity of many pesticides and there are many adjuvant formulations that have been developed for this purpose. Surfactants are nearly always components of these adjuvants ranging from minor
25 components to the sole component.

One class of adjuvants that has found success in, for example, N-phosphonomethylglycine (glyphosate) formulations containing polyoxyalkylene aliphatic amines such as, for example, ethoxylated tallowamine. While polyoxyalkylene aliphatic amine-based adjuvants have excellent surfactant
30 properties that often enhance the efficacy of phytotoxicants such as glyphosate, they unfortunately are eye irritants and must be used with a high degree of caution.

Reducing or eliminating the eye irritancy of the polyoxyalkylene aliphatic amine-based adjuvants used with pesticides, without reducing the efficacy of the

pesticidal compositions containing the surfactants, is a highly desirable end. The protection of the applicator and personnel preparing the surfactant and pesticidal compositions from eye damage is of paramount importance. Reducing the eye irritancy of the adjuvants and pesticidal compositions containing the adjuvant, increases the use that can be made of such products while lessening the possibility of injury to personnel handling and using them.

BRIEF SUMMARY OF THE INVENTION


The present invention is directed to an adjuvant composition for pesticide formulations which exhibits reduced eye irritancy. The adjuvant composition according to the invention is comprised of a polyoxyalkylene aliphatic amine containing at least about 2 moles of an alkylene oxide group and an eye irritation reducing compound which simultaneously reduces the eye irritation caused by the polyoxyalkylene aliphatic amine and which complexes with metal ions such as calcium and iron which may be present in an aqueous solution of the adjuvant and a pesticide or a plant growth regulator. The eye irritation reducing compound is a carboxylic acid that is capable of complexing with or forming chelates with metal ions in aqueous solution while simultaneously reducing the eye irritation caused by the polyoxyalkylene aliphatic amine. The adjuvants according to the invention also contain a mixture of polyhydric alcohols and, optionally, a defoamer. The mixture of polyhydric alcohols is comprised of a trihydric alcohol, such as glycerol and one or more diols, such as ethylene glycol and propylene glycol. The adjuvant according to the invention is particularly useful in N-phosphonomethylglycine (glyphosate) herbicidal formulations. The invention is also directed to a method of killing or controlling weeds which comprises contacting the weeds with a herbicidally effective amount of the composition according to the invention.

DETAILED DESCRIPTION OF THE INVENTION

The term "pesticide" as used herein includes chemicals and microbial agents used as active ingredients of products for control of crop and lawn pests and diseases, animal ectoparasites, and other pests in public health. The term also includes plant growth regulators, pest repellants, synergists, herbicide safeners (which reduce the phytotoxicity of herbicides to crop plants) and preservatives.

The adjuvant composition of the present invention comprises a polyoxyalkylene aliphatic amine having at least about 2 moles of an alkylene

group, the adjuvant composition having an eye irritancy lower than the eye irritancy of the polyoxyalkylene aliphatic amine. The polyoxyalkylene aliphatic amines according to the invention are compounds of the formula I



wherein R_1 is alkyl or alkenyl group having from 6 to 22 carbon atoms, R_2 is an alkylene group having from 2 to 4 carbon atoms, x and y are numbers such that $x + y$ has an average value of from about 2 to about 50. The polyoxyalkylene aliphatic amines are present in the adjuvant in an amount sufficient to increase the efficacy of the pesticide or plant growth regulator with which it is formulated. Such an amount of the polyoxyalkylene aliphatic amines normally causes eye irritation to users of formulations containing a pesticide or plant growth regulator, especially when such formulations are applied by spraying. Preferred polyoxyalkylene aliphatic amines are ethoxylated tallow amines having a degree of ethoxylation of about 20. In reference to compounds having the formula I, such amines are those wherein R_1 is a mixture of saturated or unsaturated carbon chains having from about 8 to about 22 carbon atoms, R_2 is an ethylene group and $x + y$ has an average value of about 20. The typical amount of polyoxyalkylene aliphatic amine in the adjuvant according to the invention can range from about 5% to about 85% by weight with the preferred amount typically ranging from about 65% to about 75% by weight.

The eye irritation reducing component of the adjuvant of the present invention is a carboxylic which is capable of forming a complex with metal ions in aqueous solution thereby reducing or eliminating the inactivating effect of metal ions on the activity of the pesticide in the formulated product. Such a carboxylic also interacts with the polyoxyalkylene aliphatic amine thereby reducing the eye irritation of the adjuvant and of any pesticide formulation containing the adjuvant.

The carboxylic acids according to the invention also function as chelating agents. It is well known that chelating agents are compounds having donor atoms that can combine by coordinate bonding with a metal ion to form a cyclic structure

known as a chelating complex. The donor atoms are present in separate functional groups within the same molecule. Thus, the carboxylic acids according to the invention are those having one or more carboxyl groups and one or more other functional groups capable of interacting with polyvalent metal ions in aqueous solution such that a stable metal chelate is formed. For example, hydroxycarboxylic acids chelate through the oxygen donor atoms located in the carboxyl group and the alcohol group. Other such carboxylic acids include, but are not limited to, aminocarboxylic acids such as ethylenediaminetetraacetic acid and its salts. N-phosphonomethylglycine (glyphosate) is an example of a herbicide that is partially or completely inactivated in aqueous solution by the presence of metal ions, particularly polyvalent metal ions such as Ca^{+2} and Fe^{+3} .

The preferred carboxylic acids are hydroxycarboxylic acids that contain one or more carboxyl groups and one or more hydroxyl groups. Such acids that are particularly useful in the practice of the present invention include, but are not limited to, citric acid, glycolic acid, gluconic acid, alpha-hydroxybutyric acid, malic acid, saccharic acid, mandelic acid, tartaric acid, glyceric acid. Citric acid is especially preferred because it is non-toxic and can be used at relatively low concentrations. Another advantage to the use of citric acid is its ability to increase the phytotoxicity of the herbicide glyphosate because citric acid readily complexes with metals such as calcium and iron, metals which are known to inactivate glyphosate. The amount of such carboxylic acids that can be present in the compositions according to the invention is an eye irritation-reducing amount which is any amount required to reduce the eye irritation of a pesticidal formulation containing the adjuvant according to the invention to an acceptable level. Such an amount will be readily determinable by those skilled in the art and will typically vary from about 0.05% to about 5% by weight of the adjuvant.

The adjuvants according to the invention also contain a mixture of polyhydric alcohols and, optionally, a defoamer. A polyhydric alcohol or polyol is a compound having at least 2 alcohol functionalities. The mixture of polyhydric alcohols is preferably comprised of at least one trihydric alcohol, preferably glycerol and at least one glycol, preferably ethylene glycol, propylene glycol or a combination thereof. While the mixture of polyhydric alcohols can contain any combination of polyols in any relative amount, it is preferably comprised of a combination of glycerol, ethylene glycol and propylene glycol present in an amount

of from about 10% to about 25% by weight of the adjuvant. The relative amounts of the various polyhydric alcohols within the mixture will vary according to the nature of the pesticide and the end use of the pesticidal formulation and will typically be ascertainable to those skilled in the art.

5 The optimum amount of each component in the adjuvant depends on variables such as the identity of the eye irritation reducing compound, the identity of the pesticide, the type of application of the pesticide composition, storage and transportation of the surfactant and pesticide compositions, the conditions of use of the pesticidal compositions, etc. and is readily determinable by those skilled in
10 the art. The adjuvant of the present invention can contain optional components to improve the water solubility of the surfactant composition and suppress gel formation. The need for such components will depend upon several factors, especially the identity of the surfactants comprising the composition.

When used as an adjuvant with glyphosate, the adjuvant according to the
15 invention is typically used at the level of from about 120 grams to about 180 grams of adjuvant per liter of an aqueous glyphosate solution containing about 480 grams of the isopropylamine salt of phosphonomethyl glycine. This is equivalent to from about 0.2% to about 0.40% by weight of a carboxylic acid having the ability to complex a metal ion.

20 The adjuvants according to the invention can be combined with a pesticidally effective amount of any type of pesticide to form a pesticidal composition. Examples of pesticides with which the adjuvants according to the invention can be formulated include, but are not limited to, glyphosate and acifluorfen (5-(2-chloro-4-(trifluoromethyl)phenoxy)-2-nitrobenzoic acid),
25 chloramben (3-amino-2,5-dichlorobenzoic acid), 2,4-D ((2,4-dichlorophenoxy)acetic acid), endotal (7-oxabicyclo(2.2.1)heptane-2,3-dicarboxylic acid), mecoprop (2-(2-methyl-4-chlorophenoxy) propionic acid), picloram (4-amino-3,5,6-trichloropyridine-2-carboxylic acid), 2,4,5-T((2,4,5-trichlorophenoxy)acetic acid), benzac (2,3,6-trichlorobenzoic acid), dicamba (3,6-dichloro-o-anisic acid), MCPA (4-chloro-o-tolyloxyacetic acid), dalapon (2,2-dichloropropionic acid), dichlorprop (2-(2,4-dichlorophenoxy)propionic acid), MCPB (4-(4-chloro-o-tolyloxy)butyric acid), bialaphos (L-2-amino-4-((hydroxy)(methyl) phosphinoyl) butyryl-L-alanyl-L-alanine), glufosinate ((3-amino-3-carboxypropyl)methylphosphinate), imazethapyr (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-

pyridinecarboxylic acid), imazaquin (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid), mixtures thereof and the like. Preferred results, however, are obtained with the herbicide glyphosate whose activity is derived from N-phosphonomethylglycine. Glyphosate is normally formulated from water soluble salts thereof. The use of glyphosate and its derivatives as herbicides is disclosed in U.S. Pat. No. 3,853,530, the entire contents of which is incorporated herein by reference. The adjuvants of the present invention are especially useful in the preparation of pesticidal compositions designed to be delivered by spraying, particularly sprayable herbicidal compositions. When combined with a pesticide, the adjuvants according to the invention can be made into a concentrate which can subsequently be diluted with water to form an aqueous pesticidal composition ready for use by spraying.

Since glyphosate in acid form has limited water solubility (about 1.2%) the water soluble salts of glyphosate are normally used for most applications. Among the water soluble salts of glyphosate are the trimethylsulfonium salt, the ammonium salt, the isopropylamine salt, and the alkali metal salts, such as sodium and potassium. These compounds due to their solubility in water are the agriculturally acceptable glyphosate-containing compounds generally used in commerce. The relative amounts of herbicide, water and surfactant in the aqueous herbicidal compositions of this invention will vary depending upon many factors including but not limited to the identity and properties of the herbicide, method of application, locus to which the herbicide is applied, etc. The weight ratio of glyphosate expressed as acid equivalent to surfactant composition is normally in the range of 1:1 to 5:1.

Formulations containing a herbicide such as glyphosate and the compositions according to the invention are effective at killing and/or controlling the growth of weeds. The following examples are meant to illustrate but not to limit the invention.

EXAMPLE 1

Adjuvant Composition

	% Weight
TRYMEEN® 6607	68.0
Citric Acid (50% Aqueous Solution)	4.0
Glycerine	15.0

AGNIQUE® DF 6889 Defoamer	0.1
Ethylene Glycol	2.0
Propylene Glycol	10.9

5 TRYMEEN® 6607 is tallow amine ethoxylated with an average of 20 moles of ethylene oxide, a trademark product of Henkel Corporation, Gulph Mills, PA 19406.

AGNIQUE® DF 6889 Defoamer is a silicone emulsion, a trademark product of Henkel Corporation, Gulph Mills, PA 19406.

EXAMPLE 2

Toxicology Testing Results

A formulation containing 480 g/L (41% by weight) of the isopropylamine salt of glyphosate and 180 g/L (15.3% by weight) of the adjuvant of Example 1 (Sample A), was compared to commercial glyphosate products under the test protocols recited below. This formulation (Sample A) contained about 0.36% citric acid by weight. Both types of ROUNDUP® contain the isopropylamine salt of glyphosate and ethoxylated tallow amine.

Table 1

SAMPLE	PDII	EYE	CLEARANCE	IRRITATION	RAINBOW TROUT LC50
A	0.2	9.0	Day 4 (S) 72 Hrs (E)	None (S) Mild (E)	35.4
B	0.7	20.0	72 Hrs (S) Day 10 (E)	Slight (S) Moderate (E)	12.9
C	0.6	19.5	72 Hrs (S) Day 7 (E)	Slight (S) Moderate (E)	5.7

Sample B is ROUNDUP®

Sample C is ROUNDUP® ULTRA

PDII is Primary Dermal Irritation Index score of the Primary Skin Irritation Test conforming to U.S. EPA Health Effects Testing Guidelines, OPPTS 870.2500 (1998) and OECD Guidelines for Testing of Chemicals, Procedure 404; and/or JMAFF 59 NohSan No. 4200, January 28, 1985.

Eye is Primary Eye Irritation Test conforming to U.S. EPA Health Effects Testing Guidelines, OPPTS 870.2500 (1998) and OECD Guidelines for Testing of Chemicals, Procedure 404; and/or JMAFF 59 Noh San No. 4200, January 28, 1985. The number is the Maximum Mean Total Score (MMTS) of this test.

Irritation- describes the level of irritation observed in the skin (S) and eye (E) tests.

Clearance- time required for irritation to clear observed in the skin (S) and eye (E) tests.

Rainbow Trout – Static Acute Toxicity Test with Rainbow Trout, *Oncorhynchus mykiss*.

Complies with U.S. EPA FIFRA Testing Guideline 72-1.

EXAMPLE 3

The post-emergence herbicidal effect of the composition of Example 2 (Sample A) was evaluated according to the following protocol:

Weed species: Barnyardgrass (ECHCG), giant foxtail (SETFA), green foxtail (SETVI), large crabgrass (DIGSA), wild oats (AVEFA), johnsongrass (SORHA), quackgrass (AGRRE), ivyleaf morningglory (IPOHE), redroot pigweed (AMARE), velvetleaf (ABUTH), cocklebur (XANST), and common lambsquarters (CHEAL), sowthistle (SONAR), Canada thistle (CIRAR), sicklepod (CASOB) and wild poinsettia (EPHHL).

Plants were germinated in synthetic soil mix (Metro-mix, manufactured by the O.M. Scott Co.) containing micronutrients under greenhouse conditions of natural light supplemented by halide light with an average energy of 165 micromols per square meter per second, daylight temperature of 29.4° C and a relative humidity of 73.7%. The photoperiod was 16 h day and 8 h night. All plants were top watered prior to treatment and sub-irrigated after treatment.

All formulations of glyphosate were brought into solution using tap water with no additional adjuvants added. The appropriate amount of compound was added to 80 ml of water to equal 800, 600, 400, 200, 100 g ae/ha when applied. Applications were made at the 2-3 leaf (B12-B13) stage with a track sprayer manufactured by Allen Machine Works utilizing a 8002E nozzle at 38 psi and a track speed of 2 mph. Application height was 18 inches above the plant canopy. This was the equivalent to an application volume of 187 L/ha. All treatments were replicated three times per species and randomized before transferring treated plant material to zone 12 or 14 for incubation during the duration of the experiment. Injury and phytotoxicity ratings were visual and taken on each pot at 7 and 14 days after treatment (DAT). The entries in the tables below range from 0% meaning no death to 100% meaning total death for a given application rate. The numbers in parentheses at the top of each table represents the number of trials conducted.

Table 2

GM/HA	IPOHE	ABUTH	SONAR	SAMPLE A		(TWO)		CASOB
				CIRAR	CHEAL	XANST	AMARE	
100	0	0	0	5.9	0	6.7	23.3	0
200	5	0.9	35	11.7	83.3	33.3	66.7	36.7
400	35	61.7	84.2	35	90	60	83.3	54.2
600	60	80.6	96.5	55	98.3	76.7	86.3	85
800	53.3	91.4	98.4	53.4	100	80	93	93.4

5

Table 3

GM/HA	EPHHL	SORHA	AVEFA	SAMPLE A		(TWO)		AGRRE
				ECHCG	DIGSA	SETVI	SETFA	
100	10	1.7	0	0	0	1.7	0	1.7
200	53.3	23.3	0.9	0	38.3	72.5	60	8.4
400	91	96.7	40	76.7	69	99.9	90	64.2
600	91.3	99.9	72.5	80	77.4	100	100	96.5
800	97.7	100	87.4	85	85.9	100	100	95

What is claimed is:

1. An adjuvant composition comprising: (a) polyoxyalkylene aliphatic amine; (b) a mixture of polyhydric alcohols; (c) an effective eye irritation reducing amount of a carboxylic acid having the ability to complex a metal ion.
- 5 2. The adjuvant of claim 1 wherein the polyoxyalkylene aliphatic amine is ethoxylated tallow amine having an average degree of ethoxylation of from about 2 to about 50.
3. The adjuvant of claim 1 wherein the average degree of ethoxylation is about 20.
- 10 4. The adjuvant of claim 1 wherein the mixture of polyhydric alcohols is comprised of glycerol, ethylene glycol and propylene glycol.
5. The adjuvant of claim 1 wherein the amount of the carboxylic acid is from about 0.05% to about 5% by weight.
- 15 6. The adjuvant of claim 1 wherein the carboxylic acid is an alpha-hydroxycarboxylic acid.
7. The adjuvant of claim 6 wherein the alpha-hydroxycarboxylic acid is citric acid.
8. The adjuvant of claim 6 wherein the amount of citric acid in the adjuvant is from about 0.05% to about 5% by weight.
- 20 9. An adjuvant composition comprising: (a) an ethoxylated tallowamine having an average degree of ethoxylation of from about 2 to about 50; (b) a mixture comprised of glycerol, ethylene glycol and propylene glycol; (c) an effective eye irritation reducing amount of citric acid.
10. The adjuvant of claim 8 wherein the average degree of ethoxylation is about 20.
- 25 11. The adjuvant of claim 8 wherein the mixture is comprised of about 15% by weight of glycerol, about 2% by weight of ethylene glycol and about 10.9% by weight of ethylene glycol.
12. The adjuvant of claim 8 wherein the amount of citric acid is about 2% by weight.
- 30 13. A pesticidal composition comprising: (a) polyoxyalkylene aliphatic amine; (b) a mixture of polyhydric alcohols; (c) an effective eye irritation reducing amount of a carboxylic acid having the ability to complex a metal ion; (d) a pesticidally effective amount of a pesticide.

14. The pesticidal composition of claim 13 wherein the polyoxyalkylene aliphatic amine is ethoxylated tallow amine having an average degree of ethoxylation of from about 2 to about 50.
- 5 15. The pesticidal composition of claim 13 wherein the average degree of ethoxylation is about 20.
16. The pesticidal composition of claim 13 wherein the mixture of polyhydric alcohols is comprised of glycerol, ethylene glycol and propylene glycol.
17. The pesticidal composition of claim 13 wherein the amount of the carboxylic acid is from about 0.05% to about 5% by weight.
- 10 18. The pesticidal composition of claim 13 wherein the carboxylic acid is an alpha-hydroxycarboxylic acid.
19. The pesticidal composition of claim 17 wherein the alpha-hydroxycarboxylic acid is citric acid.
- 15 20. The pesticidal composition of claim 17 wherein the amount of citric acid in the adjuvant is from about 0.05% to about 5% by weight.
21. A method of killing or controlling weeds comprising contacting the weeds with a herbicidally effective amount of the composition of claim 13.
- 20 22. A method of killing or controlling weeds comprising contacting the weeds with a herbicidally effective amount of a composition comprising: (a) an ethoxylated tallowamine having an average degree of ethoxylation of from about 2 to about 50; (b) a mixture comprised of glycerol, ethylene glycol and propylene glycol; (c) an effective eye irritation reducing amount of citric acid; (d) a herbicidally effective amount of N-phosphonomethylglycine or a salt thereof.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/00954

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A01N 63/00, 57/00, 25/00

US CL : 504/118, 127; 424/405

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 504/118, 127; 424/405

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y --- X	US 4,478,650 (ZADO) 23 October 1984, col. 2 lines 40-68, col. 3 lines 1-65.	1-3,5-8 ----- 1,6,7
X	GB 1081294 (DERWENT INFORMATION LTD) 28 December 1962, abstract.	1



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

&

document member of the same patent family

Date of the actual completion of the international search

21 MARCH 2000

Date of mailing of the international search report

19 MAY 2000

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Box PCT

Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ALTON PRYOR

Telephone No. (703) 308-4691